

# blood

2008 111: 5446-5456  
Prepublished online Jan 23, 2008;  
doi:10.1182/blood-2007-06-093906

## **Guidelines for the diagnosis and treatment of chronic lymphocytic leukemia: a report from the International Workshop on Chronic Lymphocytic Leukemia updating the National Cancer Institute Working Group 1996 guidelines**

Michael Hallek, Bruce D. Cheson, Daniel Catovsky, Federico Caligaris-Cappio, Guillaume Dighiero, Hartmut Döhner, Peter Hillmen, Michael J. Keating, Emili Montserrat, Kanti R. Rai and Thomas J. Kipps

---

Updated information and services can be found at:  
<http://bloodjournal.hematologylibrary.org/cgi/content/full/111/12/5446>

Articles on similar topics may be found in the following *Blood* collections:

[Neoplasia](#) (4156 articles)  
[Free Research Articles](#) (526 articles)  
[Review Articles](#) (230 articles)  
[Clinical Trials and Observations](#) (2446 articles)

---

Information about reproducing this article in parts or in its entirety may be found online at:  
[http://bloodjournal.hematologylibrary.org/misc/rights.dtl#repub\\_requests](http://bloodjournal.hematologylibrary.org/misc/rights.dtl#repub_requests)

Information about ordering reprints may be found online at:  
<http://bloodjournal.hematologylibrary.org/misc/rights.dtl#reprints>

Information about subscriptions and ASH membership may be found online at:  
<http://bloodjournal.hematologylibrary.org/subscriptions/index.dtl>

Blood (print ISSN 0006-4971, online ISSN 1528-0020), is published semimonthly by the American Society of Hematology, 1900 M St, NW, Suite 200, Washington DC 20036.  
[Copyright 2007 by The American Society of Hematology; all rights reserved.](#)



## Review article

# Guidelines for the diagnosis and treatment of chronic lymphocytic leukemia: a report from the International Workshop on Chronic Lymphocytic Leukemia updating the National Cancer Institute–Working Group 1996 guidelines

Michael Hallek,<sup>1</sup> Bruce D. Cheson,<sup>2</sup> Daniel Catovsky,<sup>3</sup> Federico Caligaris-Cappio,<sup>4</sup> Guillaume Dighiero,<sup>5</sup> Hartmut Döhner,<sup>6</sup> Peter Hillmen,<sup>7</sup> Michael J. Keating,<sup>8</sup> Emili Montserrat,<sup>9</sup> Kanti R. Rai,<sup>10</sup> and Thomas J. Kipps<sup>11</sup>

<sup>1</sup>Klinik I für Innere Medizin, Universität zu Köln, Köln, Germany; <sup>2</sup>Georgetown University Hospital, Lombardi Cancer Center, Washington, DC; <sup>3</sup>Institute of Cancer Research, London, United Kingdom; <sup>4</sup>Università Vita-Salute Istituto di Ricovero e Cura a Carattere Scientifico (IRCSS) San Raffaele, Milano, Italy; <sup>5</sup>Institute Pasteur, Montevideo, Uruguay; <sup>6</sup>University of Ulm, Ulm, Germany; <sup>7</sup>Pinderfields Hospital, Wakefield, United Kingdom; <sup>8</sup>Department of Leukemia, University of Texas, M. D. Anderson Cancer Center, Houston; <sup>9</sup>Hospital Clinic, Barcelona, Spain; <sup>10</sup>Division of Hematology/Oncology, Long Island Jewish Medical Center, New Hyde Park, NY; and <sup>11</sup>Rebecca and John Moores Cancer Center, University of California–San Diego, La Jolla

**Standardized criteria for diagnosis and response assessment are needed to interpret and compare clinical trials and for approval of new therapeutic agents by regulatory agencies. Therefore, a National Cancer Institute–sponsored Working Group (NCI-WG) on chronic lymphocytic leukemia (CLL) published**

**guidelines for the design and conduct of clinical trials for patients with CLL in 1988, which were updated in 1996. During the past decade, considerable progress has been achieved in defining new prognostic markers, diagnostic parameters, and treatment options. This prompted the International Workshop on Chronic Lym-**

**phocytic Leukemia (IWCLL) to provide updated recommendations for the management of CLL in clinical trials and general practice. (Blood. 2008;111:5446-5456)**

© 2008 by The American Society of Hematology

## Introduction

In 1988 and 1996, a National Cancer Institute-sponsored Working Group (NCI-WG) on chronic lymphocytic leukemia (CLL) published guidelines for the design and conduct of clinical trials for patients with CLL to facilitate comparisons between different treatments and to establish definitions that could be used in scientific studies on the biology of this disease.<sup>1,2</sup> The Food and Drug Administration also adopted these guidelines in their evaluation and approval of new drugs. During the last decade, considerable progress has been made in defining new prognostic markers, diagnostic parameters, and treatment options, prompting the IWCLL-sponsored Working Group to revise the 1996 criteria.

## Diagnosis of CLL

The World Health Organization classification of hematopoietic neoplasias describes CLL as leukemic, lymphocytic lymphoma, being only distinguishable from small lymphocytic lymphoma (SLL) by its leukemic appearance.<sup>3</sup> In the World Health Organization classification, CLL is always a disease of neoplastic B cells, whereas the entity formerly described as T-CLL is now called T-cell prolymphocytic leukemia.<sup>4</sup>

It is important to verify that the patient has CLL and not some other lymphoproliferative disease that can masquerade as CLL, such as hairy cell leukemia, or leukemic manifestations of mantle cell lymphoma, marginal zone lymphoma, splenic marginal zone lymphoma with circulating villous lymphocytes, or follicular lymphoma. To achieve this, it is essential to evaluate the blood

count, blood smear, and the immune phenotype of the circulating lymphoid cells (see “Blood” and “Immunophenotype”).

## Blood

The diagnosis of CLL requires the presence of more than or equal to  $5 \times 10^9/L$  B lymphocytes ( $5000/\mu L$ ) in the peripheral blood for the duration of at least 3 months. The clonality of the circulating B lymphocytes needs to be confirmed by flow cytometry. The leukemia cells found in the blood smear are characteristically small, mature lymphocytes with a narrow border of cytoplasm and a dense nucleus lacking discernible nucleoli and having partially aggregated chromatin. These cells may be found admixed with larger or atypical cells, cleaved cells, or prolymphocytes, which may comprise up to 55% of the blood lymphocytes.<sup>5</sup> Finding prolymphocytes in excess of this percentage would favor a diagnosis of prolymphocytic leukemia (B-cell PLL). Gumprecht nuclear shadows, or smudge cells, found as cell debris, are other characteristic morphologic features found in CLL.

CLL or SLL might be suspected in otherwise healthy adults who have an absolute increase in the clonal B lymphocytes but who have less than  $5 \times 10^9/L$  B lymphocytes in the blood. However, in the absence of lymphadenopathy or organomegaly (as defined by physical examination and CT scans), cytopenias, or disease-related symptoms, the presence of fewer than  $5 \times 10^9/L$  B lymphocytes blood is defined as “monoclonal B-lymphocytosis.”<sup>6</sup> The presence of a cytopenia caused by a typical marrow infiltrate defines the diagnosis of CLL regardless of the number of peripheral blood B lymphocytes or of the lymph node involvement. monoclonal B-lymphocytosis seems to progress to frank CLL at a rate

Submitted June 2, 2007; accepted December 15, 2007. Prepublished online as *Blood* First Edition paper, January 23, 2008; DOI 10.1182/blood-2007-06-093906.

The online version of this article contains a data supplement.

© 2008 by The American Society of Hematology

of 1% to 2% per year (A. C. Rawstron, F. L. Bennett, M. O'Connor, P. H., manuscript submitted, May 2007).

The definition of SLL requires the presence of lymphadenopathy and the absence of cytopenias caused by a clonal marrow infiltrate. Moreover, the number of B lymphocytes in the peripheral blood should not exceed  $5 \times 10^9/L$ . In SLL, the diagnosis should be confirmed by histopathologic evaluation of a lymph node biopsy whenever possible.

### Immunophenotype

CLL cells coexpress the T-cell antigen CD5 and B-cell surface antigens CD19, CD20, and CD23. The levels of surface immunoglobulin, CD20, and CD79b are characteristically low compared with those found on normal B cells.<sup>7,8</sup> Each clone of leukemia cells is restricted to expression of either kappa or lambda immunoglobulin light chains.<sup>7</sup> Variations of the intensity of expression of these markers may exist and do not prevent inclusion of a patient in clinical trials for CLL.

In contrast, B-cell PLL cells do not express CD5 in half of the cases, and typically express high levels of CD20 and surface Ig.<sup>9</sup> In addition, the leukemia cells of mantle cell lymphoma, despite also expressing B-cell surface antigens and CD5, generally do not express CD23.

### Other tests performed at diagnosis

The tests described in this section are not needed to establish the diagnosis of CLL but may help predict the prognosis or to assess the tumor burden. With the exception of molecular genetics fluorescence in situ hybridization (FISH), the application of these tests should not be used in routine practice to influence therapy and is not generally recommended. However, certain parameters, such as immunoglobulin mutational status, are useful for predicting the clinical course in individual cases. These tests can be recommended for patients who want a better prediction of the rate at which their disease might progress but it should be emphasized that the indication for treatment does not depend on any of these tests but on the clinical stage and the disease activity (see "Indications for treatment").

**Molecular genetics.** Using interphase FISH, cytogenetic lesions can be identified in more than 80% of all CLL cases.<sup>10</sup> The most common deletions are in the long arm of chromosome 13 [del(13q14.1)]. Additional, frequent chromosomal aberrations comprise deletions and/or trisomy of chromosome 12, deletions in the long arm of chromosomes 11 [del(11q)] and 6 [del(6q)], and in the short arm of chromosome 17 [del(17p)].<sup>10</sup> When stimulated in vitro, CLL cells can have detectable chromosomal translocations, which are of potential prognostic significance.<sup>11</sup> However, certain translocations can help distinguish other lymphoproliferative diseases from CLL [eg, t(11;14)], which generally is found in mantle cell lymphoma).

There is increasing evidence from prospective clinical trials that detection of certain chromosomal deletions has prognostic significance. Patients with leukemia cells that have del(17p) have an inferior prognosis and appear resistant to standard chemotherapy regimens using alkylating drugs and/or purine analogs.<sup>12,13</sup> In a retrospective analysis on several chromosomal aberrations as detected by FISH, patients who had CLL cells with chromosomal aberrations del(11q) and del(17p) had an inferior outcome compared with patients who had leukemia cells with a normal karyotype or del(13q) as the sole genetic abnormality.<sup>10</sup> On the other hand, patients with leukemia cells

having del(17p) may respond to therapy with alemtuzumab, either alone or in combination with other antileukemia agents.<sup>14,15</sup> Detection of these cytogenetic abnormalities has apparent prognostic value and may influence therapeutic decisions. For clinical trials, it is recommended that cytogenetics be performed before treating a patient on protocol. Additional genetic defects may be acquired during the course of the disease<sup>16</sup>; therefore, the repetition of FISH analyses seems justified before subsequent, second- and third-line treatment.

**Mutational status of IgV<sub>H</sub>, VH3.21 usage, and expression of ZAP-70 or CD38.** The leukemia cells express immunoglobulin that may or may not have incurred somatic mutations in the immunoglobulin heavy chain variable region genes (IgV<sub>H</sub> genes). The outcome of patients with leukemia cells that use an unmutated IgV<sub>H</sub> gene is inferior to those patients with leukemia cells that use a mutated IgV<sub>H</sub> gene.<sup>17,18</sup> In addition, the VH3.21 gene usage is an unfavorable prognostic marker independent of the IgV<sub>H</sub> mutational status.<sup>19</sup> Leukemia-cell expression of ZAP-70 and CD38 was found to correlate with the expression of unmutated IgV<sub>H</sub> genes and to predict a poor prognosis.<sup>20-26</sup> However, the association between expression of ZAP-70 or CD38 with the expression of unmutated IgV<sub>H</sub> genes is not absolute. It is uncertain whether leukemia-cell expression of unmutated IgV<sub>H</sub> genes or ZAP-70 predict the response to treatment or overall survival, once therapy is required.<sup>13,27</sup> Taken together, further clinical trials are needed to standardize the assessment of these parameters and to determine whether they should affect the management of patients with CLL.

**Serum markers.** Several studies have found that serum markers CD23, thymidine kinase, and  $\beta_2$ -microglobulin may predict survival or progression-free survival.<sup>28-34</sup> Assays for these markers should be standardized and used in prospective clinical trials to validate their relative value to the management of patients with CLL.

**Marrow examination.** In CLL, characteristically more than 30% of the nucleated cells in the aspirate are lymphoid. Although the type of marrow infiltration (diffuse vs. nondiffuse) reflects the tumor burden and provides some prognostic information, recent studies of the German and Spanish study groups suggest that the prognostic value of BM biopsy may now be superseded by new prognostic markers.

A marrow aspirate and biopsy generally are not required for the diagnosis of CLL. However, a marrow biopsy and aspirate can help evaluate for factors that might contribute to cytopenias (anemia, thrombocytopenia) that may or may not be directly related to leukemia-cell infiltration of the marrow. Because such factors could influence the susceptibility to drug-induced cytopenias, a marrow biopsy is recommended before initiating therapy. It is recommended to repeat a marrow biopsy in patients with persisting cytopenia after treatment to uncover disease- versus therapy-related causes.

## Clinical staging

There are 2 widely accepted staging methods for use in both patient care and clinical trials: the Rai<sup>35</sup> and the Binet<sup>36</sup> systems. The original Rai classification was modified to reduce the number of prognostic groups from 5 to 3.<sup>37</sup> As such, both systems now describe 3 major subgroups with discrete clinical outcomes. These 2 staging systems are simple, inexpensive, and can be applied by physicians worldwide. Both solely rely on a physical examination

and standard laboratory tests and do not require ultrasound, computed tomography (CT), or magnetic resonance imaging. These 2 systems are outlined in the following 2 sections.

### Rai staging system

The modified Rai classification defines low-risk disease as patients who have lymphocytosis with leukemia cells in the blood and/or marrow (lymphoid cells > 30%; formerly considered Rai stage 0). Patients with lymphocytosis, enlarged nodes in any site, and splenomegaly and/or hepatomegaly (lymph nodes being palpable or not) are defined as having intermediate risk disease (formerly considered Rai stage I or stage II). High-risk disease includes patients with disease-related anemia (as defined by a hemoglobin [Hb] level <110 g/L [11 g/dL]; formerly stage III) or thrombocytopenia (as defined by a platelet count <100 × 10<sup>9</sup>/L; formerly stage IV).

### Binet staging system

Staging is based on the number of involved areas, as defined by the presence of enlarged lymph nodes of greater than 1 cm in diameter or organomegaly, and on whether there is anemia or thrombocytopenia.

The areas of involvement considered for staging are as follows: (1) Head and neck, including the Waldeyer ring (this counts as one area, even if more than one group of nodes is enlarged). (2) Axillae (involvement of both axillae counts as one area). (3) Groins, including superficial femorals (involvement of both groins counts as one area). (4) Palpable spleen. (5) Palpable liver (clinically enlarged).

**Stage A.** Hb 100 g/L (10 g/dL) or more and platelets 100 × 10<sup>9</sup>/L or more and up to 2 of the above areas involved.

**Stage B.** Hb 10 g/L or more and platelets 100 × 10<sup>9</sup>/L or more and organomegaly greater than that defined for stage A (ie, 3 or more areas of nodal or organ enlargement).

**Stage C.** All patients who have Hb less than 10 g/L and/or a platelet count less than 100 × 10<sup>9</sup>/L, irrespective of organomegaly.

## Eligibility criteria for clinical trials

The selection of CLL patients for clinical trials is similar to that for patients with other malignancies. Phase 1 or 2 clinical trials commonly, although not invariably, are intended for patients who have had prior therapy. It may be worth considering the inclusion of patients with SLL in some phase 1 or 2 trials exploring new agents in CLL. However, for SLL the response assessment should be done according to the lymphoma guidelines. The combination of new agents with standard therapy as part of phase 2 studies may be investigated in both untreated and previously treated patients. Phase 3 clinical trials are used to compare the clinical outcome using new treatment modalities with that using current standard therapy. Other requirements for eligibility with respect to age, clinical stage, performance status, organ function, or status of disease activity should be defined for each study.

### Performance status and fitness

Before inclusion in a trial, the performance status as defined by the Eastern Cooperative Oncology Group should be 0 to 3. Future clinical trials involving elderly patients ideally should assess the comorbidity (fitness) and/or functional activity of patients (eg, such

as that defined by “cumulative illness rating scale” or the “Charlson” score).<sup>38,39</sup>

### Organ function eligibility for clinical trials

Most chemotherapy agents have potential toxicity for the liver, kidneys, heart, lungs, nervous system, or other organ systems. Therefore, organ function requirements should be guided by the known or suspected toxicity of each agent based on preclinical studies or prior clinical studies. Patients enrolled on protocols evaluating agents with known or suspected toxicity for a given organ(s) should have documented the specific organ function before therapy.

### Infectious disease status

The status of specific infectious diseases as outlined in “Required pretreatment evaluation” should be documented. Patients with active infections requiring systemic antibiotics, antifungal or antiviral drugs should have their infection resolved before initiating therapy in a clinical trial.

### Second malignancies

Patients with a second malignancy, other than basal cell carcinoma of the skin or in situ carcinoma of the cervix or the breast, generally are not considered candidates for entry into clinical trials unless the tumor was successfully treated with curative intent at least 2 years before trial entry.

### Required pretreatment evaluation

Parameters considered necessary for a complete pretreatment evaluation may differ depending on whether or not the patient is treated in a clinical protocol. Therefore, a clear distinction is made in this section and in “Definition of response, relapse, and refractory disease” between recommendations for general practice and the requirements for clinical trials (Tables 1-3). If not indicated otherwise, recommendations are identical for clinical trials and general practice. In general, studies for defining these parameters should be performed within 2 weeks of clinical trial enrollment (except for marrow aspirate and biopsy and CT scans (see “Essential pretreatment tests”).

#### Essential pretreatment tests.

**Physical examination.** The bidimensional diameters of the largest palpable lymph nodes in each of the following sites should be recorded: cervical, axillary, supraclavicular, inguinal, and femoral (Table 1). The size of the liver and spleen, as assessed by palpation, should also be recorded.

#### Assessment of performance status.

**Complete blood cell count.** White blood cell count, hemoglobin and hematocrit, platelet count, and differential count, including both percent and absolute number of lymphocytes, and reticulocyte count should be performed. Reporting the proportion of prolymphocytes is desirable when these are present.

**Marrow biopsy.** Before initiating treatment in a clinical trial with potentially myelosuppressive agents, patients should undergo a unilateral marrow aspirate and biopsy. Repeat marrow biopsies may be compared with the pretreatment marrow specimen.

**Serum chemistry.** (eg, creatinine, bilirubin, lactic dehydrogenase, transaminases, alkaline phosphatase).

#### Serum immunoglobulin levels.

#### Direct antiglobulin test.

**Chest radiograph.** (When a CT scan is not performed.)

**Table 1. Pretreatment evaluation of patients with CLL**

Diagnostic test	General practice*	Clinical trial
<b>Tests to establish the diagnosis†</b>	—	—
Complete blood count and differential count‡	Always	Always
Immunophenotyping of lymphocytes§	Always	Always
<b>Assessment before treatment¶</b>	—	—
History and physical, performance status	Always	Always
Complete blood count and differential**	Always	Always
Marrow aspirate and biopsy	Desirable	Desirable
Serum chemistry, serum immunoglobulin, direct antiglobulin test‡‡	Always	Always
Chest radiograph§§	Always	Always
Infectious disease status¶¶	Always	Always
<b>Additional tests before treatment   </b>	—	—
Cytogenetics (FISH) for del(13q), del(11q), del(17p), and (12), del(6q) in the peripheral blood lymphocytes***	Desirable	Always
IgVH mutational status, ZAP-70, and CD38§	No	Always
CT scan of chest, abdomen, and pelvis†††	No	Desirable
MRI, lymphangiogram, gallium scan, PET scans‡‡‡	No	No
Abdominal ultrasound§§§	Possible	No

The section of guidelines for each table entry is indicated by symbols.

No indicates not generally indicated; RQ, research question; PBS, peripheral blood smear; MRI, magnetic resonance imaging; PET, positron emission tomography; FISH, fluorescence in situ hybridization.

\*General practice is defined as the use of accepted treatment options for a patient with CLL not enrolled in a clinical trial.

†“Diagnosis of CLL.”

‡“Blood.”

§“Immunophenotype.”

¶“Essential pretreatment tests.”

||“Physical examination,” “Assessment of performance status.”

\*\*“Complete blood cell count.”

‡‡“Marrow biopsy.”

‡‡‡“Serum chemistry,” “Serum immunoglobulin levels,” “Direct antiglobulin test.”

§§“Chest radiograph.”

¶¶“Infectious disease status.”

|||“Additional pretreatment tests (Table 1) may be performed in clinical trials or in the presence of specific clinical problems.”

\*\*\*“The assessment of molecular cytogenetics (FISH) before therapy is recommended.”

†††“CT scans generally are not required for the initial evaluation or follow-up.”

‡‡‡“Other imaging methods.”

§§§“Abdominal ultrasound.”

**HIV.** Patients who are infected with HIV should be given special consideration because of the potential risks for immune suppression with most antileukemia therapies and the potential for compounded myelotoxicity of treatment with antiretroviral therapy.

**Cytomegalovirus (CMV).** Therapies associated with the potential for reactivation of infection with CMV, such as alemtuzumab or allogeneic stem cell transplantation, should include plans for monitoring for active CMV disease and/or for providing anti-CMV therapy.<sup>40</sup> These should cover screening or early diagnosis of CMV reactivation and its subsequent management. However, a positive CMV serology does not represent a contraindication for alemtuzumab treatment or allogeneic stem cell transplantation. As a general recommendation for patients treated with alemtuzumab, close monitoring and/or therapy for active CMV disease should be considered for patients found to have evidence for increased levels of CMV in the blood by the polymerase chain reaction (PCR), even in the absence of clinical symptoms. In addition, evaluation and therapy for CMV is recommended for any patient with clinical symptoms of active CMV infection.

**Hepatitis B and hepatitis C.** Before initiating treatment, the evaluation for infection with hepatitis B virus (HBV) and hepatitis C virus (HCV) is recommended because reactivation of HBV and HCV infections may occur under therapy with immunosuppressive or myelosuppressive drugs. Chronic HBV carriers as defined by positive surface antigen undergoing chemotherapy should receive prophylactic therapy with nucleoside analogs, such as lamivudine to prevent HBV reactivation.<sup>41,42</sup>

**Additional pretreatment tests (Table 1) may be performed in clinical trials or in the presence of specific clinical problems.**

*The assessment of molecular cytogenetics (FISH) before therapy is recommended.*

*CT scans generally are not required for the initial evaluation or follow-up.* Moreover, the staging of CLL does not use CT scans but relies on physical examination and blood counts. A recent study has found that patients in Rai stage 0 but with detectable abdominal disease by CT scans may have a more aggressive disease.<sup>43</sup> Therefore, clinical studies evaluating the use of CT scans in CLL are strongly encouraged. Moreover, enlarged lymph nodes if detected only by CT scan do not change the clinical Binet or Rai stage.

In clinical trials where the treatment intent is to maximize complete remission (CR), chest, abdominal, and pelvic CT scans are recommended to evaluate the response to therapy. Two CT scans should be performed, one before the start of therapy and one at the first restaging after therapy if previously abnormal.

**Other imaging methods.** Except in some patients with Richter transformation, positron emission tomography scans do not provide information that is useful in the management of CLL. Similarly, nuclear magnetic resonance imaging and other imaging techniques are generally not useful in the management of CLL.

**Abdominal ultrasound.** In some countries, the use of abdominal ultrasound is popular to assess the extent of lymphadenopathy and organomegaly in CLL. Although it may be used in the clinical management of individual patients, this methodology is strongly

**Table 2. Recommendations regarding indications for treatment in CLL**

	General practice*	Clinical trial
Treat with Rai stage 0	No†	RQ
Treat with Binet stage A	No†	RQ
Treat with Binet stage B or Rai stage I or II	Possible†	Possible†
Treat with Binet stage C or Rai stage III or IV	Yes	Yes
Treatment of active/progressive disease	Yes	Yes
Treat without active/progressive disease	No	RQ

No indicates not generally indicated; RQ, research question.

\*General practice is defined as the use of accepted treatment options for a patient with CLL not enrolled in a clinical trial.

†Treatment is indicated, if the disease is active as defined in "Indications for treatment."

investigator-dependent and should therefore not be used for the response evaluation in clinical trials.

A lymph node biopsy is generally not required, unless such tissue is necessary for companion scientific studies or in rare cases with difficult diagnosis. A lymph node biopsy is requested to establish the diagnosis of a transformation into an aggressive lymphoma (Richter syndrome).

## Indications for treatment

### Primary treatment decisions

Criteria for initiating treatment may vary depending on whether or not the patient is treated in a clinical trial (Table 2). In general practice, newly diagnosed patients with asymptomatic early-stage disease (Rai 0, Binet A) should be monitored without therapy unless they have evidence of disease progression. Studies from both the French Cooperative Group on CLL,<sup>44</sup> the Cancer and Leukemia Group B,<sup>45</sup> the Spanish Group Pethema,<sup>46</sup> and the Medical Research Council<sup>47</sup> in the United Kingdom in patients with early-stage disease confirm that the use of alkylating agents in patients with such early-stage disease does not prolong survival. This result was confirmed by a meta-analysis.<sup>47</sup> In one study, treated patients with early-stage disease had an increased frequency of fatal epithelial cancers compared with untreated patients.<sup>44</sup> Therefore, the potential benefit, if any, of an early-intervention therapy with antileukemia drugs, alone or in combination with monoclonal antibodies, requires further study.

Whereas patients at intermediate (stage I and II) and high risk (stage III and IV) according to the modified Rai classification or at Binet stage B or C usually benefit from the initiation of treatment, some of these patients (in particular Rai intermediate risk or Binet stage B) can be monitored without therapy until they have evidence for progressive or symptomatic disease.

Active disease should be clearly documented for protocol therapy. At least one of the following criteria should be met: (1) Evidence of progressive marrow failure as manifested by the development of, or worsening of, anemia and/or thrombocytopenia. (2) Massive (ie, > 6 cm below the left costal margin) or progressive or symptomatic splenomegaly. (3) Massive nodes (ie, > 10 cm in longest diameter) or progressive or symptomatic lymphadenopathy. (4) Progressive lymphocytosis with an increase of more than 50% over a 2-month period or lymphocyte doubling time (LDT) of less than 6 months. LDT can be obtained by linear regression extrapolation of absolute lymphocyte counts obtained at intervals of 2 weeks over an observation period of 2 to 3 months; patients with initial blood lymphocyte counts of less than  $30 \times 10^9/L$

( $30\,000/\mu L$ ) may require a longer observation period to determine the LDT. In addition, factors contributing to lymphocytosis or lymphadenopathy other than CLL (eg, infections) should be excluded. (5) Autoimmune anemia and/or thrombocytopenia poorly responsive to corticosteroids or other standard therapy (see "Autoimmune hemolytic anemia or autoimmune thrombocytopenia"). (6) A minimum of any one of the following disease-related symptoms must be present: (a) Unintentional weight loss more than or equal to 10% within the previous 6 months. (b) Significant fatigue (ie, Eastern Cooperative Oncology Group PS 2 or worse; cannot work or unable to perform usual activities). (c) Fevers of greater than  $100.5^\circ F$  or  $38.0^\circ C$  for 2 or more weeks without other evidence of infection. (d) Night sweats for more than 1 month without evidence of infection.

Hypogammaglobulinemia or monoclonal or oligoclonal paraproteinemia does not by itself constitute a basis for initiating therapy. However, it is recommended to assess the change of these protein abnormalities if patients are treated.

Patients with CLL may present with a markedly elevated leukocyte count; however, the symptoms associated with leukocyte aggregates that develop in patients with acute leukemia rarely occur in patients with CLL. Therefore, the absolute lymphocyte count should not be used as the sole indicator for treatment

### Second-line treatment decisions

In general, second-line treatment decisions follow the same indications as those used for initiation of first-line treatment. Patients who have resistant disease, a short time to progression after the first treatment, and/or leukemia cells with del(17p) often do not respond to standard chemotherapy and have a relatively short survival. Therefore, such patients should be offered investigative clinical protocols, including allogeneic hematopoietic stem cell transplantation.<sup>48-52</sup>

## Definition of response, relapse, and refractory disease

Assessment of response should include a careful physical examination and evaluation of the blood and marrow. Imaging studies, in particular CT scans, generally are not required except to monitor the response to therapy in clinical trials (Table 3).

### Complete remission

CR requires all of the following criteria as assessed at least 3 months after completion of therapy:

**Absence of clonal lymphocytes in the peripheral blood.**

**Absence of significant lymphadenopathy.** (eg, lymph nodes > 1.5 cm in diameter) by physical examination. In clinical trials, a CT scan of the abdomen, pelvis, and thorax is desirable if previously abnormal. Lymph nodes should not be larger than 1.5 cm in diameter.

**No hepatomegaly or splenomegaly by physical examination.** In clinical trials, a CT scan of the abdomen should be performed at response assessment if found to be abnormal before therapy or if physical examination is inconclusive at the time of evaluation.

**Absence of constitutional symptoms.**

**Blood counts above the following values.** (1) Polymorphonuclear leukocytes  $1.5 \times 10^9/L$  ( $1500/\mu L$ ) or more. (2) Platelets more than  $100 \times 10^9/L$  ( $100\,000/\mu L$ ). (3) Hemoglobin more than 110 g/L ( $11.0\text{ g/dL}$ ; untransfused).

**Table 3. Recommendations regarding the response assessment in CLL patients**

Diagnostic test	General practice*	Clinical trial
History, physical examination†	Always	Always
Immunophenotyping of peripheral blood lymphocytes‡	If clinical and hematologic response indicates CR	If clinical response and hematologic response indicates CR
CBC and differential count§	Always	Always
Marrow aspirate and biopsy¶	At cytopenia of uncertain cause	At CR or cytopenia of uncertain cause
Assessment for minimal residual disease	No	If a long-lasting CR is the desired endpoint
Ultrasound of the abdomen†	Possible, if previously abnormal	No
CT scans of chest, pelvis, and abdomen†	No	Indicated if previously abnormal and otherwise in CR

The section of guidelines for each table entry is indicated by symbols.

\*General practice is defined as the use of accepted treatment options for a patient with CLL not enrolled in a clinical trial.

†Absence of significant lymphadenopathy; ‡No hepatomegaly or splenomegaly by physical examination; §Reduction in lymphadenopathy; ¶Absence of clonal lymphocytes in the peripheral blood.

‡Absence of clonal lymphocytes in the peripheral blood.

§Blood counts above the following values: †The blood count should show one of the following results, †An increase in the number of blood lymphocytes by 50% or more with at least 5000 B lymphocytes per microliter, †Occurrence of cytopenia (neutropenia, anemia, or thrombocytopenia) attributable to CLL.

¶For patients in clinical trials (Table 3): a marrow aspirate and biopsy should be performed at least 3 months after the last treatment and if clinical and laboratory results (the first 5 points under "Complete remission") demonstrate that a CR has been achieved.

||Minimal residual disease.

**For patients in clinical trials (Table 3); a marrow aspirate and biopsy should be performed at least 3 months after the last treatment and if clinical and laboratory results (the first 5 points under "Complete remission") demonstrate that a CR has been achieved.** The marrow should be analyzed by flow cytometry and/or immunohistochemistry to demonstrate that the marrow is free of clonal B-CLL cells. Cases with residual CLL cells by conventional (not 4-color; see "Minimal residual disease") flow cytometry or immunohistochemistry are defined as partial remission (PR).

In some cases, lymphoid nodules can be found (formerly used to define nodular PR), which often reflect residual disease.<sup>53,54</sup> Therefore, these nodules should be assessed by immunohistochemistry to define whether they are comprised primarily of T cells or lymphocytes other than CLL cells or of CLL cells. The category of "nodular PR" should no longer be used. If the marrow is found to be hypocellular, a repeat marrow biopsy should be performed after 4 to 6 weeks, provided that the blood counts have recovered. Marrow biopsies should be compared with that of the pretreatment marrow. In some cases, it is necessary to postpone the marrow biopsy until after all the other criteria to define a CR have been satisfied. However, this time interval should not exceed 6 months after the last treatment.

**A controversial issue is how best to categorize the response of patients who fulfill all the criteria for a CR (including the marrow examinations) but who have a persistent anemia or thrombocytopenia or neutropenia apparently unrelated to CLL but related to drug toxicity.** We recommend that these patients should be considered as a different category of remission, CR with incomplete bone marrow recovery (CRi). For the definition of this category, CRi, the marrow evaluation should be performed with scrutiny and not show any clonal infiltrate. In clinical trials, CRi patients should be monitored prospectively to determine whether their outcome differs from that of patients with detectable residual disease or with noncytopenic CR.

### Partial remission

PR is defined by the criteria described in the 2 following paragraphs, and/or the third paragraph below (if abnormal before therapy), as well as one or more of the features listed in the fourth paragraph below. To define a PR, at least one of these parameters needs to be documented for a minimal duration of 2 months (Table 4). Constitutional symptoms persisting for more than 1 month should be recorded.

**A decrease in the number of blood lymphocytes by less than 50% or more from the value before therapy.**

**Reduction in lymphadenopathy.** To be assessed by CT scans in clinical trials<sup>55</sup> or by palpation in general practice as defined by the following: (1) A decreased lymph node size by 50% or more in the sum products of up to 6 lymph nodes, or in one lymph node diameter if only a single lymph node was present before therapy. (2) No increase in any lymph node, and no new enlarged lymph node. In small lymph nodes (< 2 cm), an increase of less than 25% is not considered to be significant.

**A decrease in the size of the liver and/or spleen by 50% or more as defined by CT scan in clinical trials or by palpation or ultrasound in general practice.**

**The blood count should show one of the following results:**

(1) Polymorphonuclear leukocytes at  $1.5 \times 10^9/L$  ( $1500/\mu L$ ) or more or 50% improvement over baseline without granulocyte-colony-stimulating factor (G-CSF) support. (2) Platelet counts greater than  $100 \times 10^9/L$  ( $100\,000/\mu L$ ) or 50% improvement over baseline. (3) Hemoglobin greater than 110 g/L (11.0 g/dL) or 50% improvement over baseline without red blood cell transfusions or erythropoietin support.

### Progressive disease

Progressive disease is characterized by at least one of the following:

**Lymphadenopathy.** Progression of lymphadenopathy is discovered almost uniformly by blood counts. Therefore, imaging methods are not needed to follow CLL progression. Progression of lymphadenopathy may be documented by physical examination. Disease progression occurs if one of the following events is observed: (1) Appearance of any new lesion, such as enlarged lymph nodes (> 1.5 cm), splenomegaly, hepatomegaly or other organ infiltrates. (2) An increase by 50% or more in greatest determined diameter of any previous site. A lymph node of 1 to 1.5 cm must increase by 50% or more to a size greater than 1.5 cm in the longest axis. A lymph node of more than 1.5 cm must increase to more than 2.0 cm in the longest axis. (3) An increase of 50% or more in the sum of the product of diameters of multiple nodes.

**An increase in the liver or spleen size by 50% or more or the de novo appearance of hepatomegaly or splenomegaly.**

**An increase in the number of blood lymphocytes by 50% or more with at least 5000 B lymphocytes per microliter.**

**Table 4. Response definition after treatment for patients with CLL, using the parameters of Tables 1 and 3**

Parameter	CR	PR	PD	SD
<b>Group A</b>				
Lymphadenopathy*	None more than 1.5 cm	Decrease $\geq$ 50%	Increase $\geq$ 50%	Change of -49% to +49%
Liver and/or spleen size	Normal size	Decrease $\geq$ 50%	Increase $\geq$ 50%	Change of -49% to +49%
Constitutional symptoms	None	Any	Any	Any
Polymorphonuclear leukocytes	>1500/ $\mu$ L	>1500/ $\mu$ L or > 50% improvement over baseline	Any	Any
Circulating clonal B lymphocytes	None	Decrease $\geq$ 50% from baseline	Increase $\geq$ 50% over baseline	Change of -49% to +49%
<b>Group B</b>				
Platelet count	>100 000/ $\mu$ L	>100 000/ $\mu$ L or increase $\geq$ 50% over baseline	Decrease of $\geq$ 50% from baseline secondary to CLL	Change of -49 to +49%
Hemoglobin	>11.0 g/dL (untransfused and without erythropoietin)	>11 g/dL or increase $\geq$ 50% over baseline	Decrease of >2 g/dL from baseline secondary to CLL	Increase $\leq$ 11.0 g/dL or <50% over baseline, or decrease <2 g/dL
Marrow	Normocellular, <30% lymphocytes, no B-lymphoid nodules; hypocellular marrow defines CRi	$\geq$ 30% lymphocytes, or B-lymphoid nodules, or not done	Increase of lymphocytes to more than 30% from normal	No change in marrow infiltrate

Values in SI units for leukocytes are  $1.5 \times 10^9/L$ ; for platelets  $100 \times 10^9/L$ ; and for hemoglobin 110 g/L and 20 g/L.

CR indicates complete remission (all of the criteria have to be met); PR, partial remission (at least one of the criteria of group A and one of the criteria of group B have to be met); PD, progressive disease (at least one of the criteria of group A or one of the criteria of group B have to be met); SD, stable disease (all of the above criteria have to be met).

\* Sum of the products of multiple lymph nodes (as evaluated by CT scans in clinical trials, or by physical examination or ultrasound in general practice).

**Transformation to a more aggressive histology (eg, Richter syndrome).** Whenever possible, this diagnosis should be established by lymph node biopsy.

**Occurrence of cytopenia (neutropenia, anemia, or thrombocytopenia) attributable to CLL.** (1) During therapy: cytopenias may occur as a side effect of many therapies and should be assessed according to Table 5. During therapy, cytopenias cannot be used to define disease progression. Each protocol should define the amount of drug(s) to be administered with such cytopenias. (2) After treatment: The progression of any cytopenia (unrelated to autoimmune cytopenia), as documented by a decrease of Hb levels by more than 20 g/L (2 g/dL) or to less than 100 g/L (10 g/dL), or by a decrease of platelet counts by more than 50% or to less than  $100 \times 10^9/L$  ( $100\,000/\mu L$ ), which occurs at least 3 months after

treatment, defines disease progression, if the marrow biopsy demonstrates an infiltrate of clonal CLL cells.

#### Stable disease

Patients who have not achieved a CR or a PR, and who have not exhibited progressive disease, will be considered to have stable disease (which is equivalent to a nonresponse).

#### Beneficial responses

Responses that should be considered clinically beneficial include CR and PR; all others (eg, stable disease, nonresponse, progressive disease, or death from any cause) should be rated as a treatment failure.

**Table 5. Grading scale for hematologic toxicity in CLL studies**

Grade*	Decrease in platelets† or Hb‡ (nadir) from pretreatment value, %	Absolute neutrophil count/ $\mu$ L§ (nadir)
0	No change to 10%	$\geq$ 2000
1	11%-24%	$\geq$ 1500 and <2000
2	25%-49%	$\geq$ 1000 and <1500
3	50%-74%	$\geq$ 500 and <1000
4	$\geq$ 75%	<500

\*Grades: 1, mild; 2, moderate; 3, severe; 4, life-threatening; 5, fatal. Death occurring as a result of toxicity at any level of decrease from pretreatment will be recorded as grade 5.

†Platelet counts must be below normal levels for grades 1 to 4. If, at any level of decrease, the platelet count is less than  $20 \times 10^9/L$  ( $20\,000/\mu L$ ), this will be considered grade 4 toxicity, unless a severe or life-threatening decrease in the initial platelet count (eg,  $20 \times 10^9/L$ ) was present pretreatment, in which case the patient is not evaluable for toxicity referable to platelet counts.

‡Hb levels must be below normal levels for grades 1 to 4. Baseline and subsequent Hb determinations must be performed before any given transfusions. The use of erythropoietin is irrelevant for the grading of toxicity but should be documented.

§If the absolute neutrophil count (ANC) reaches less than  $10^9/L$  ( $1000/\mu L$ ), it should be judged to be grade 3 toxicity. Other decreases in the white blood cell count, or in circulating granulocytes, are not to be considered because a decrease in the white blood cell count is a desired therapeutic endpoint. A gradual decrease in granulocytes is not a reliable index in CLL for stepwise grading of toxicity. If the ANC was less than  $10^9/L$  before therapy, the patient is not evaluable for toxicity referable to the ANC. The use of G-CSF is irrelevant for the grading of toxicity but should be documented.

#### Duration of response and progression-free survival

Duration of response should be measured from the end of the last treatment until evidence of progressive disease (as defined above). Progression-free survival is defined as the interval between the first treatment day to the first sign of disease progression. Event-free survival is defined as the interval between the first treatment day to the first sign of disease progression, or treatment for relapse, or death (whichever occurs first). Survival duration is defined as the interval between the first treatment day to death.

#### Relapse

Relapse is defined as a patient who has previously achieved the above criteria ("Complete remission," "Partial remission") of a CR or PR, but after a period of 6 or more months, demonstrate evidence of disease progression (see preceding discussion of progressive disease).

#### Refractory disease

Refractory disease is defined as treatment failure (as defined in "Beneficial responses") or disease progression within 6 months to the last antileukemic therapy. For the definition of "high risk CLL" justifying the use of allogeneic stem cell transplantation,<sup>56</sup> the

disease should be refractory to a purine-analog based therapy or to autologous hematopoietic stem cell transplantation.

### Minimal residual disease

The complete eradication of the leukemia is an obvious desired end point. New detection technologies, such as multicolor flow cytometry and real-time quantitative PCR, have determined that many patients who achieved a complete response by the 1996 NCI-WG guidelines have detectable minimal residual disease (MRD). Although eradication of MRD may improve prognosis, prospective clinical trials are needed to define whether additional treatment intended solely to eradicate MRD provides a significant benefit to clinical outcome. The techniques for assessing MRD have undergone a critical evaluation and have become fairly standard.<sup>57</sup> Either 4-color flow cytometry (MRD flow) or allele-specific oligonucleotide PCR is reliably sensitive down to a level of approximately one CLL cell in 10 000 leukocytes. As such, patients will be defined as having a clinical remission in the absence of MRD when they have blood or marrow with less than one CLL cell per 10 000 leukocytes. The blood generally can be used for making this assessment except during the period within 3 months of completing therapy, particularly for patients treated with alemtuzumab, rituximab, and other antibodies targeting CLL. In such cases, it is essential to assess the marrow for MRD. Therefore, future clinical trials that aim toward achieving long-lasting CRs should include at least one test to assess MRD because the lack of leukemia persistence using these sensitive tests seems to have a strong, positive prognostic impact.<sup>58-60</sup>

### Factors requiring stratification at inclusion in a clinical phase 3 trial

Patients ideally should be stratified with regard to previous treatment versus no previous treatment, and as purine analog-sensitive versus purine-analog refractory in studies for which prior therapy is allowed. If more than one clinical stage is allowed, patients ideally should be stratified for stage. Patients ideally should be stratified based upon whether or not they have leukemia cells with del(17p) or del(11q).

### Assessment of toxicity

Evaluation of treatment-related toxicity requires careful consideration of both the manifestations of the underlying disease and the anticipated adverse reactions to the agents used in therapy. For this reason, some of the conventional criteria used for assessing toxicity are not applicable to clinical studies involving patients with hematologic malignancies in general, or CLL in particular. An example is hematologic toxicity; patients with advanced CLL generally have cytopenias that may be caused by the underlying CLL and/or prior therapy. A few guidelines are presented to help evaluate for treatment-induced toxicity in CLL.

#### Hematologic toxicity

Evaluation of hematologic toxicity in patients with CLL must take into consideration that many patients have low blood cell counts at the initiation of therapy. Therefore, the standard criteria used for solid tumors cannot be applied, as many CLL patients then would

be considered to have grade 2 to 4 hematologic toxicity at the initiation of treatment. Furthermore, the absolute blood neutrophil counts rarely are used at the initiation of therapy to modify the treatment dose because these values typically are unreliable in CLL patients with lymphocytosis. However, the increasing use of more effective therapeutic agents, particularly those with neutropenia as a dose-limiting toxicity (eg, nucleoside analogs), can result in clinically significant myelosuppression. Therefore, the 1996 guidelines proposed a new dose-modification scheme for quantifying hematologic deterioration in patients with CLL, which included alterations in the dose of myelosuppressive agents based on the absolute neutrophil count. This dose modification scheme has proven very helpful in the context of several large prospective trials in CLL and should be retained (Table 5).

#### Infectious complications

Patients with CLL are at increased risk for infection because of compromised immune function, which might be related to the disease itself and/or to the consequences of therapy. Nevertheless, the rate(s) of infection after treatment can be used in assessing the relative immune-suppressive effects of a given therapy. The etiology of the infection should be reported and categorized as bacterial, viral, or fungal, and as proven or probable. The severity of infections should be quantified as minor (requiring either oral antimicrobial therapy or symptomatic care alone), major (requiring hospitalization and systemic antimicrobial therapy), or fatal (death as a result of the infection).

Particular attention should be given to monitoring for symptoms or laboratory evidence of infection with CMV in patients treated with agents, such as alemtuzumab (alone or in combination) or with allogeneic stem cell transplantation. In contrast, the infection rate seems low in patients younger than 65 years treated with fludarabine-based first-line therapy, where no monitoring or routine anti-infective prophylaxis is required.<sup>61</sup>

#### Tumor lysis syndrome

CLL patients rarely experience tumor lysis syndrome after therapy with a purine analog-based regimen.<sup>62</sup> However, this might not be the case after treatment with newer agents or novel treatment modalities. For this reason, patients in early-phase clinical trials should be monitored for possible tumor lysis syndrome, which should be treated appropriately. If observed, the occurrence and severity of tumor lysis syndrome should be recorded in clinical trials.

#### Nonhematologic toxicities

Other nonhematologic toxicities should be graded according to the latest version of the NCI Common Toxicity Criteria.<sup>63</sup>

### Reporting clinical response data

Clear and careful data reporting is an essential part of any clinical trial. In clinical studies involving previously treated patients, patients who are relapsed or refractory should be clearly distinguished. Relapse and refractory disease are defined in "Relapse" and "Refractory disease." For those patients who have relapsed, it is also useful to describe the quality and duration of their prior response.

## Treatment endpoints

Given the recent increase of treatment options for CLL patients, the choice of treatment and the endpoints of clinical trials may depend on the fitness of the patients (see "Performance status and fitness"). For example, the number of MRD-negative CRs or the overall survival might be appropriate endpoints in physically fit patients. In contrast, trials on patients with reduced physical fitness might choose the time to progression or health-related quality of life as trial endpoints. Moreover, recent data suggest that the quality of life in CLL patients is reduced compared with the normal population and only moderately increased by some of the current treatment options.<sup>64-67</sup> Therefore, further studies assessing the health-related quality of life in CLL are strongly encouraged.

## Supportive care and management of complications

### Indications for growth factors in CLL

While under myelosuppressive (chemo-)therapy, growth factors, such as G-CSF, should be given according to the guidelines of the American Society of Clinical Oncology.<sup>68</sup> The use of G-CSF also might benefit patients who experience prolonged cytopenias after treatment with alemtuzumab. Similarly, some CLL patients with anemia may benefit from erythropoiesis stimulating factors if used according to recently published guidelines.<sup>69,70</sup> However, it should be pointed out that CLL-related cytopenias are often efficiently corrected by an appropriate antileukemic therapy.

### Autoimmune hemolytic anemia or autoimmune thrombocytopenia

Immune thrombocytopenic purpura (ITP) and autoimmune hemolytic anemia (AIHA) as a single abnormality caused by CLL initially should be treated with glucocorticoids and not chemotherapy. Second-line treatment options for AIHA include splenectomy, intravenous immunoglobulins, and/or immunosuppressive therapy with agents, such as cyclosporine A, azathioprine, or low-dose cyclophosphamide. Good responses also have been obtained with antibody therapy using agents as rituximab or alemtuzumab.<sup>71-73</sup> Treatment refractory autoimmune cytopenias can be an indication for chemotherapy or chemoimmunotherapy directed at the underlying CLL.<sup>74</sup> In this regard, the Binet or Rai

staging systems do not distinguish between ITP/AIHA or marrow infiltration as the cause for anemia or thrombocytopenia that results in classifying a patient as having stage C or high-risk disease.

## Acknowledgments

The authors thank the following colleagues for reading and commenting on the current version of the guidelines: Drs Francesc Bosch (Barcelona, Spain), John Byrd (Columbus, OH), Barbara Eichhorst (Cologne, Germany), Terry Hamblin (London, United Kingdom), Neil Kay (Rochester, MN), Eva Kimby (Stockholm, Sweden), Estella Matutes (London, United Kingdom), Stefano Molica (Catanzaro, Italy), Stephen Mulligan (Sydney, Australia), Susan O'Brien (Houston, TX), David Oscier (Bournemouth, United Kingdom), and John Seymour (Melbourne, Australia). The authors also thank the following colleagues for their active participation in the original version of the guidelines: Charles A. Schiffer, Martin M. Oken, David H. Boldt, Sanford J. Kempin, and Kenneth A. Foon. The authors also thank the following colleagues for their active participation of the second, 1996 version of the guidelines: John M. Bennett, Michael Grever, Neil Kay, and Susan O'Brien.

This work was supported in part by the Competence Network Malignant Lymphoma of the German Ministry for Education and Research as well as German Cancer Aid (Deutsche Krebshilfe; M.H.), the European Leukemia Network (European Research Initiative on CLL, ERIC; M.H., H.D.), of the National Institutes of Health PO1-CA081534 to the CLL Research Consortium, the Arbib Foundation (D.C.), the Karches Family Foundation, Prince Family Foundation, and the Nash Family Foundation (K.R.R.), and Associazione Italiana per la Ricerca sul Cancro (AIRC; F.C.-C.).

## Authorship

Contribution: M.H., B.D.C., D.C., F.C.-C., G.D., H.D., P.H., M.J.K., E.M., K.R.R., and T.J.K. all wrote the paper.

A complete list of the International Workshop on Chronic Lymphocytic Leukemia is provided on the *Blood* website; see the Supplemental Materials link at the top of the online article.

Conflict-of-interest disclosure: The authors declare no competing financial interests.

Correspondence: Michael Hallek, Klinik I für Innere Medizin, Universität zu Köln, Joseph-Stelzmann Str. 9, 50924 Köln, Germany; e-mail: michael.hallek@uni-koeln.de.

## References

- Cheson BD, Bennett JM, Rai KR, et al. Guidelines for clinical protocols for chronic lymphocytic leukemia (CLL): recommendations of the NCI-sponsored working group. *Am J Hematol*. 1988; 29:153-163.
- Cheson BD, Bennett JM, Grever M, et al. National Cancer Institute-Sponsored Working Group guidelines for chronic lymphocytic leukemia: revised guidelines for diagnosis and treatment. *Blood*. 1996;87:4990-4997.
- Müller-Hermelink HK, Montserrat E, Catovsky D, Harris NL. Chronic lymphocytic leukemia/small lymphocytic lymphoma. In: Jaffe ES, Harris NL, Stein H, Vardiman JW, eds. *World Health Organization Classification of Tumours: Pathology and Genetics of Tumours of Haematopoietic and Lymphoid Tissues*. Lyon, France: IARC Press; 2001: 127-130.
- Catovsky D, Ralfkiaer E, Müller-Hermelink HK. T-cell prolymphocytic leukaemia. In: Jaffe ES, Harris NL, Stein H, Vardiman JW, eds. *World Health Organization Classification of Tumours: Pathology and Genetics of Tumours of Haematopoietic and Lymphoid Tissues*. Lyon, France: IARC Press; 2001:195-196.
- Melo JV, Catovsky D, Galton DAG. The relationship between chronic lymphocytic leukaemia and prolymphocytic leukaemia: IV. Analysis of survival and prognostic features. *Br J Haematol*. 1986;63: 377-387.
- Marti GE, Rawstron AC, Ghia P, et al. Diagnostic criteria for monoclonal B-cell lymphocytosis. *Br J Haematol*. 2005;130:325-332.
- Moreau EJ, Matutes E, A'Hern RP, et al. Improvement of the chronic lymphocytic leukemia scoring system with the monoclonal antibody SN8 (CD79b). *Am J Clin Pathol*. 1997;108:378-382.
- Ginaldi L, De Martinis M, Matutes E, Farahat N, Morilla R, Catovsky D. Levels of expression of CD19 and CD20 in chronic B cell leukaemias. *J Clin Pathol*. 1998;51:364-369.
- Catovsky D, Müller-Hermelink HK, Montserrat E, Harris NL. B-cell prolymphocytic leukaemia. In: Jaffe ES, Harris NL, Stein H, Vardiman JW, eds. *World Health Organization Classification of Tumours: Pathology and Genetics of Tumours of Haematopoietic and Lymphoid Tissues*. Lyon, France: IARC Press; 2001:131-132.
- Döhner H, Stilgenbauer S, Benner A, et al. Genomic aberrations and survival in chronic lymphocytic leukemia. *N Engl J Med*. 2000;343: 1910-1916.
- Mayr C, Speicher MR, Kofler DM, et al. Chromosomal translocations are associated with poor prognosis in chronic lymphocytic leukemia. *Blood*. 2006;107:742-751.
- Döhner H, Fischer K, Bentz M, et al. p53 gene

- deletion predicts for poor survival and nonresponse to therapy with purine analogs in chronic B-cell leukemias. *Blood*. 1995;85:1580-1589.
13. Grever MR, Lucas DM, Dewald GW, et al. Comprehensive assessment of genetic and molecular features predicting outcome in patients with chronic lymphocytic leukemia: results from the US Intergroup Phase III Trial E2997. *J Clin Oncol*. 2007;25:799-804.
  14. Stilgenbauer S, Döhner H. Campath-1H-induced complete remission of chronic lymphocytic leukemia despite p53 gene mutation and resistance to chemotherapy. *N Engl J Med*. 2002;347:452-453.
  15. Lozanski G, Heerema NA, Finni IW, et al. Alemtuzumab is an effective therapy for chronic lymphocytic leukemia with p53 mutations and deletions. *Blood*. 2004;103:3278-3281.
  16. Shanafelt TD, Witzig TE, Fink SR, et al. Prospective evaluation of clonal evolution during long-term follow-up of patients with untreated early-stage chronic lymphocytic leukemia. *J Clin Oncol*. 2006;24:4634-4641.
  17. Damle RN, Wasil T, Fais F, et al. Ig V gene mutation status and CD38 expression as novel prognostic indicators in chronic lymphocytic leukemia [see comments]. *Blood*. 1999;94:1840-1847.
  18. Hamblin TJ, Davis Z, Gardiner A, Oscier DG, Stevenson FK. Unmutated Ig V(H) genes are associated with a more aggressive form of chronic lymphocytic leukemia [see comments]. *Blood*. 1999;94:1848-1854.
  19. Thorselius M, Krober A, Murray F, et al. Strikingly homologous immunoglobulin gene rearrangements and poor outcome in VH3-21-using chronic lymphocytic leukemia patients independent of geographic origin and mutational status. *Blood*. 2006;107:2889-2894.
  20. Orchard JA, Ibbotson RE, Davis Z, et al. ZAP-70 expression and prognosis in chronic lymphocytic leukaemia. *Lancet*. 2004;363:105-111.
  21. Crespo M, Bosch F, Villamor N, et al. ZAP-70 expression as a surrogate for immunoglobulin-variable-region mutations in chronic lymphocytic leukemia. *N Engl J Med*. 2003;348:1764-1775.
  22. Rassenti LZ, Huynh L, Toy TL, et al. ZAP-70 compared with immunoglobulin heavy-chain gene mutation status as a predictor of disease progression in chronic lymphocytic leukemia. *N Engl J Med*. 2004;351:893-901.
  23. Ibrahim S, Keating MJ, Do KA, et al. CD38 expression as an important prognostic factor in B-cell chronic lymphocytic leukemia. *Blood*. 2001;98:181-186.
  24. Lin K, Sherrington PD, Dennis M, Matrai Z, Cawley JC, Pettitt AR. Relationship between p53 dysfunction, CD38 expression, and IgV(H) mutation in chronic lymphocytic leukemia. *Blood*. 2002;100:1404-1409.
  25. Ghia P, Guida G, Stella S, et al. The pattern of CD38 expression defines a distinct subset of chronic lymphocytic leukemia (CLL) patients at risk of disease progression. *Blood*. 2003;101:1262-1269.
  26. Boonstra JG, van Lom K, Langerak AW, et al. CD38 as a prognostic factor in B cell chronic lymphocytic leukaemia (B-CLL): comparison of three approaches to analyze its expression. *Cytometry B Clin Cytom*. 2006;70:136-141.
  27. Byrd JC, Gribben JG, Peterson BL, et al. Select high-risk genetic features predict earlier progression following chemoimmunotherapy with fludarabine and rituximab in chronic lymphocytic leukemia: justification for risk-adapted therapy. *J Clin Oncol*. 2006;24:437-443.
  28. Hallek M, Langenmayer I, Nerl C, et al. Elevated serum thymidine kinase levels identify a subgroup at high risk of disease-progression in early, non-smoldering chronic lymphocytic leukemia. *Blood*. 1999;93:1732-1737.
  29. Keating MJ, Lerner S, Kantarjian H, Freireich EJ, O'Brien S. The serum  $\beta$ 2-microglobulin ( $\beta$ 2m) level is more powerful than stage in predicting response and survival in chronic lymphocytic leukemia (CLL). *Blood*. 1995;86(Suppl 1):606a.
  30. Reinisch W, Wilhelm M, Hilgarth M, et al. Soluble CD23 reliably reflects disease activity in B-cell chronic lymphocytic leukemia. *J Clin Oncol*. 1994;12:2146-2149.
  31. Sarfati M, Chevret S, Chastang C, et al. Prognostic importance of serum soluble CD23 level in chronic lymphocytic leukemia. *Blood*. 1996;88:4259-4264.
  32. Wierda WG, O'Brien S, Wang X, et al. Prognostic nomogram and index for overall survival in previously untreated patients with chronic lymphocytic leukemia. *Blood*. 2007;109:4679-4685.
  33. Magnac C, Porcher R, Davi F, et al. Predictive value of serum thymidine kinase level for Ig-V mutational status in B-CLL. *Leukemia*. 2003;17:133-137.
  34. Matthews C, Catherwood MA, Morris TC, et al. Serum TK levels in CLL identify Binet stage A patients within biologically defined prognostic subgroups most likely to undergo disease progression. *Eur J Haematol*. 2006;77:309-317.
  35. Rai KR, Sawitsky A, Cronkite EP, Chanana AD, Levy RN, Pasternack BS. Clinical staging of chronic lymphocytic leukemia. *Blood*. 1975;46:219-234.
  36. Binet JL, Auquier A, Dighiero G, et al. A new prognostic classification of chronic lymphocytic leukemia derived from a multivariate survival analysis. *Cancer*. 1981;48:198-204.
  37. Rai KR. A critical analysis of staging in CLL. In: Gale RP, Rai KR, eds. *Chronic Lymphocytic Leukemia: Recent Progress and Future Directions*. New York, NY: Liss; 1987:253-264.
  38. Extermann M, Overcash J, Lyman GH, Parr J, Balducci L. Comorbidity and functional status are independent in older patients. *J Clin Oncol*. 1998;16:1582-1587.
  39. Balducci L, Extermann M. Management of cancer in the older person: a practical approach. *Oncologist*. 2000;5:224-237.
  40. O'Brien SM, Keating MJ, Mocarski ES. Updated guidelines on the management of cytomegalovirus reactivation in patients with chronic lymphocytic leukemia treated with alemtuzumab. *Clin Lymphoma Myeloma*. 2006;7:125-130.
  41. Yagci M, Acar K, Sucak GT, Aki Z, Bozdayi G, Haznedar R. A prospective study on chemotherapy-induced hepatitis B virus reactivation in chronic HBs Ag carriers with hematologic malignancies and pre-emptive therapy with nucleoside analogues. *Leuk Lymphoma*. 2006;47:1608-1612.
  42. Rossi G, Pelizzari A, Motta M, Puoti M. Primary prophylaxis with lamivudine of hepatitis B virus reactivation in chronic HBsAg carriers with lymphoid malignancies treated with chemotherapy. *Br J Haematol*. 2001;115:58-62.
  43. Muntanola A, Bosch F, Arguis P, et al. Abdominal computed tomography predicts progression in patients with Rai stage 0 chronic lymphocytic leukemia. *J Clin Oncol*. 2007;25:1576-1580.
  44. Dighiero G, Maloum K, Desablens B, et al. Chlorambucil in indolent chronic lymphocytic leukemia. *N Engl J Med*. 1998;338:1506-1514.
  45. Shustik C, Mick R, Silver R, Sawitsky A, Rai K, Shapiro L. Treatment of early chronic lymphocytic leukemia: intermittent chlorambucil versus observation. *Hematol Oncol*. 1988;6:7-12.
  46. Montserrat E, Fontanillas M, Estape J, for the Spanish PETHEMA Group. Chronic lymphocytic leukemia treatment: an interim report of PET-HEMA trials. *Leuk Lymphoma*. 1991;5:89-92.
  47. CLL Trialists' Collaborative Group. Chemotherapeutic options in chronic lymphocytic leukemia: a meta-analysis of the randomized trials. *J Natl Cancer Inst*. 1999;91:861-868.
  48. Moreno C, Villamor N, Colomer D, et al. Allogeneic stem-cell transplantation may overcome the adverse prognosis of unmutated VH gene in patients with chronic lymphocytic leukemia. *J Clin Oncol*. 2005;23:3433-3438.
  49. Dreger P, Brand R, Milligan D, et al. Reduced-intensity conditioning lowers treatment-related mortality of allogeneic stem cell transplantation for chronic lymphocytic leukemia: a population-matched analysis. *Leukemia*. 2005;19:1029-1033.
  50. Gribben JG, Zahrieh D, Stephans K, et al. Autologous and allogeneic stem cell transplantations for poor-risk chronic lymphocytic leukemia. *Blood*. 2005;106:4389-4396.
  51. Schetelig J, Thiede C, Bornhauser M, et al. Evidence of a graft-versus-leukemia effect in chronic lymphocytic leukemia after reduced-intensity conditioning and allogeneic stem-cell transplantation: the Cooperative German Transplant Study Group. *J Clin Oncol*. 2003;21:2747-2753.
  52. Caballero D, Garcia-Marco JA, Martino R, et al. Allogeneic transplant with reduced intensity conditioning regimens may overcome the poor prognosis of B-cell chronic lymphocytic leukemia with unmutated immunoglobulin variable heavy-chain gene and chromosomal abnormalities (11q- and 17p-). *Clin Cancer Res*. 2005;11:7757-7763.
  53. Oudat R, Keating MJ, Lerner S, O'Brien S, Albitar M. Significance of the levels of bone marrow lymphoid infiltrate in chronic lymphocytic leukemia patients with nodular partial remission. *Leukemia*. 2002;16:632-635.
  54. Noy A, Verma R, Glenn M, et al. Clonotypic polymerase chain reaction confirms minimal residual disease in CLL nodular PR: results from a sequential treatment CLL protocol. *Blood*. 2001;97:1929-1936.
  55. Cheson BD, Pfistner B, Juweid ME, et al. Revised response criteria for malignant lymphoma. *J Clin Oncol*. 2007;25:579-586.
  56. Dreger P, Corradini P, Kimby E, et al. Indications for allogeneic stem cell transplantation in chronic lymphocytic leukemia: the EBMT transplant consensus. *Leukemia*. 2007;21:12-17.
  57. Rawstron AC, Villamor N, Ritgen M, et al. International standardized approach for flow cytometric residual disease monitoring in chronic lymphocytic leukaemia. *Leukemia*. 2007;21:956-964.
  58. Moreton P, Kennedy B, Lucas G, et al. Eradication of minimal residual disease in B-cell chronic lymphocytic leukemia after alemtuzumab therapy is associated with prolonged survival. *J Clin Oncol*. 2005;23:2971-2979.
  59. Wendtner CM, Ritgen M, Schweighofer CD, et al. Consolidation with alemtuzumab in patients with chronic lymphocytic leukemia (CLL) in first remission: experience on safety and efficacy within a randomized multicenter phase III trial of the German CLL Study Group (GCLLSG). *Leukemia*. 2004;18:1093-1101.
  60. Bosch F, Ferrer A, Lopez-Guillermo A, et al. Fludarabine, cyclophosphamide and mitoxantrone in the treatment of resistant or relapsed chronic lymphocytic leukaemia. *Br J Haematol*. 2002;119:976-984.
  61. Eichhorst BF, Busch R, Schweighofer C, Wendtner CM, Emmerich B, Hallek M. Due to low infection rates no routine anti-infective prophylaxis is required in younger patients with chronic lymphocytic leukaemia during fludarabine-based first line therapy. *Br J Haematol*. 2007;136:63-72.
  62. Cheson BD, Frame JN, Vena D, Quashu N, Soresen JM. Tumor lysis syndrome: an uncommon complication of fludarabine therapy of chronic lymphocytic leukemia. *J Clin Oncol*. 1998;16:2313-2320.
  63. National Cancer Institute. Common Toxicity Criteria. <http://ctep.cancer.gov/reporting/ctc.html>. Accessed April 15, 2008.
  64. Eichhorst BF, Busch R, Obwandner T, Kuhn-Hallek I, Hershbach P, Hallek M. Health-related

- quality of life in younger patients with chronic lymphocytic leukemia treated with fludarabine plus cyclophosphamide or fludarabine alone for first-line therapy: a study by the German CLL Study Group. *J Clin Oncol*. 2007;25:1722-1731.
65. Molica S. Quality of life in chronic lymphocytic leukemia: a neglected issue. *Leuk Lymphoma*. 2005;46:1709-1714.
66. Holzner B, Kemmler G, Kopp M, Nguyen-Van-Tam D, Sperner-Unterwieser B, Greil R. Quality of life of patients with chronic lymphocytic leukemia: results of a longitudinal investigation over 1 yr. *Eur J Haematol*. 2004;72:381-389.
67. Levy V, Porcher R, Delabarre F, Leporrier M, Cazin B, Chevret S. Evaluating treatment strategies in chronic lymphocytic leukemia: use of quality-adjusted survival analysis. *J Clin Epidemiol*. 2001;54:747-754.
68. Ozer H, Armitage JO, Bennett CL, et al. 2000 update of recommendations for the use of hematopoietic colony-stimulating factors: evidence-based, clinical practice guidelines. American Society of Clinical Oncology Growth Factors Expert Panel. *J Clin Oncol*. 2000;18:3558-3585.
69. Ludwig H, Rai K, Blade J, et al. Management of disease-related anemia in patients with multiple myeloma or chronic lymphocytic leukemia: epoetin treatment recommendations. *Hematol J*. 2002;3:121-130.
70. Lichtin A. The ASH/ASCO clinical guidelines on the use of erythropoietin. *Best Pract Res Clin Haematol*. 2005;18:433-438.
71. Rodon P, Breton P, Courouble G. Treatment of pure red cell aplasia and autoimmune haemolytic anaemia in chronic lymphocytic leukaemia with Campath-1H. *Eur J Haematol*. 2003;70:319-321.
72. Hamblin TJ. Autoimmune complications of chronic lymphocytic leukemia. *Semin Oncol*. 2006;33:230-239.
73. Zaja F, Vianelli N, Sperotto A, et al. Anti-CD20 therapy for chronic lymphocytic leukemia-associated autoimmune diseases. *Leuk Lymphoma*. 2003;44:1951-1955.
74. Gupta N, Kavuru S, Patel D, et al. Rituximab-based chemotherapy for steroid-refractory autoimmune hemolytic anemia of chronic lymphocytic leukemia. *Leukemia*. 2002;16:2092-2095.